

FloroPhos®

DIRECTIONS FOR USE

- After thorough prophylaxis, fill trays to approximately 1/3 capacity. Reusable or disposable trays may be used. Airdry the teeth and insert trays for 1 - 4 minutes (after 1 minute, the material is approximately 80% effective). Instruct patient to bite down lightly with intermittent chewing motion to enhance interproximal coverage.
- 2. Remove the trays and instruct patient to expectorate any excess gel. Patient should not eat, drink or rinse mouth for at least thirty minutes.

Alternatively, the gel may be applied with plastic cups or cotton applicators.

STORAGE AND SHELF-LIFE

The shelf-life of *FloroPhos* is two years when stored at temperatures between $15^{\circ}F(-10^{\circ})$ and $83^{\circ}F(30^{\circ})$.

CAUTION: U.S. FEDERAL LAW PROHIBITS DISPENSING WITHOUT PRESCRIPTION

Manufactured by Scientific Pharmaceuticals, Inc. 3221 Producer Way, Pomona, CA, U.S.A. 91768 Phone: (800) 634-3047 • (909) 595-9922

Net 32 fl.oz. (0.95L)

CONTAINS 1.23% FLUORIDE ION DERIVED FROM SODIUM FLUORIDE AND HYDROFLUORIC ACID IN A BUFFERED 0.1 MOLAR PHOSPHORIC ACID SOLUTION.

For Professional Dental Use Only